



THE INTERNATIONAL MOUNTAINEERING AND CLIMBING FEDERATION
UNION INTERNATIONALE DES ASSOCIATIONS D'ALPINISME

Office: Monbijoustrasse 61 • Postfach
CH-3000 Berne 23 • SWITZERLAND
Tel.: +41 (0)31 3701828 • Fax: +41 (0)31 3701838
e-mail: office@uiaa.ch

CONSENSUS STATEMENT

OF THE

UIAA MEDICAL COMMISSION

VOL: 15

Work in Hypoxic Conditions

Intended for Physicians, Occupational Health and Safety
Advisors, Interested Non-medical Persons

Th. Küpper, J.S. Milledge, D. Hillebrandt, J. Kubalova, U. Hefti, B. Bas-
nayt, U. Gieseler, V. Schöffl

2009

1. Introduction

Apart from “classical” hypoxia, which is almost an exclusive problem for mountaineers and, to lesser extent, for travelling business men, hypoxia is now also being used for fire prevention in storage areas, for altitude training in sport and for many other purposes. As a consequence more and more employees and other persons are exposed to hypoxia. Up to now there has been no consensus on how to provide occupational health and safety advice. Most regulations do not take into account the kind of exposure or other circumstances (e.g. whether the person is able to “escape” from hypoxia at any time). Regulations do not define the type or degree of different risks – if any – and therefore a more specialised analysis of the individual exposure is necessary to provide adequate advice for health and safety. As the world’s umbrella organization for activity in a low oxygen environment, the Medical Commission of the Union Internationale des Associations d’Alpinisme (UIAA) has a special responsibility to coordinate an international consensus on this topic. For easier use the paper is structured as follows:

- Differentiation between several types of exposure to hypoxic conditions and the consequences of each one
- Procedures of preventive medical care in Occupational Medicine and Safety for people working in the different hypoxic situations
- Comments on the differentiation and procedures

Although there are some minor physiological differences between simulated (isobaric) and “real” altitude (hypobaric hypoxia, altitude sojourn), these are not relevant for occupational safety and health. Therefore the term “altitude” includes in this paper the situation of “simulated altitude” (achieved by isobaric hypoxia) or “equivalent altitude” (a term, which is often used for aircraft cabin pressure).

Facilities with isobaric hypoxia can be left immediately and it is easy possible to survey employees working in such facilities much better (“controlled hypoxia”). Therefore a significant lower risk can be assumed in such facilities than during real altitude sojourns where it is neither possible to “escape” from hypoxia within a short time nor is it possible to perform a good survey of the health status of the person by independent (non-hypoxic) persons (“uncontrolled hypoxia”).

Note: This paper focuses on altitude related aspects of Occupational Medicine only! Other aspects of Occupational Medicine are excluded here and have to be taken into account for the individual activity at altitude. An example for this will be given in appendix 1 for persons with high workload at high altitude, such as when working in alpine (helicopter) rescue organizations.

2. Types of occupational exposure to hypoxic conditions

With regard to the current discussion in several countries (Austria, G.B., Finland, Germany) it should be emphasised that (mild) hypoxia is not normally a risk. Five important factors must be taken into account for the differentiation and the risk profile of exposure to hypoxia:

- Altitude or equivalent altitude (% O₂), respectively

- Duration of exposure
- Altitude profile / acclimatization (intermittent hypoxia included)
- Workload in hypoxic condition
- Native Highlanders vs. native lowlanders

Individual risks caused by pre-existing conditions will be discussed later (see also ref [1]).

With the five major points given above at least four kinds of exposure each with a completely different risk profile can be established (fig.1).




Figure 1: Different types of occupational exposure to hypoxic conditions (see also table 1)

2.1 Extreme Short Exposure

Extreme short exposure normally takes place at altitudes between **1800m and 2500m** and for a period for some **minutes to some hours**. With isobaric hypoxia of about 17.0-14.8% O₂ (+/- 0.2) in rooms equipped with hypoxia systems for fire prevention, expose employees to an altitude of about 1,700-2,600m (based on ICAO standard atmosphere, fig.2). This altitude is well within the range of the so-called “threshold altitude”, which is the altitude, where the body shows the first responses to hypoxia. Depending on the system observed, the threshold altitude varies between 1,500m (slight increase in pulse rate at rest) to 2,400m (increase of erythropoietin serum concentration) [2], [3]. In conclusion altitudes around the threshold do not cause any risk for healthy people by hypoxia. It also does not cause risks for people with chronic diseases of moderate severity [1]. Severely ill persons will be discussed later.

In some special circumstances there are some employees exposed to equivalent altitudes between 2,700m and 3,800m in rooms for fire protection. This situation is limited in the duration of exposure to a maximum of some hours, often for <60 min. As for any room equipped with hypoxia systems for fire protection these employees can leave immediately at any time if they do not feel well.

Some common activities expose people to even higher altitude, e.g. skiing to 3,800m (Europe) or >4,000m (U.S.A), or road traffic (nearly 3,000m (Europe), >4,000m (U.S.A, & Tibet), >5,000m (South America). Here the main problem may be the change of pressure, especially for children or persons suffering from an infection of the upper airways.

The longest exposure of this type (“extreme short exposure”) is long distance flights which can be classed as “limited exposure cases”. There are data that some airlines operate even higher cabin altitude than the limit of 2,400m given by ICAO, especially when using modern aircraft [4]. Normally the exposure time does not exceed a few hours. Up to 3,000m (and even higher) there is no risk to develop any altitude disease during this period [5], [6], [7], [8]. The main problem for this group may be the acute change of pressure, especially in the case if a person suffers from a common cold. Normally everybody – including pregnant women [9], [10] and children [11] – will feel well at these altitudes. Exceptions are people with severe pre-existing diseases (see specific paragraph below and table 3).

Group	Typical (equivalent) altitude	Typical duration of exposure	Typical Risk
Extreme short exposure (aircrafts, cable cars, skiing, road traffic, fire protection rooms)	1,800 – 2,600m	minutes to hours	Pressure changes (persons with infection of the upper airways)
	2,600 – 3,800m		No risk by hypoxia for any person who does not have severe cardiopulmonary illness No risk for healthy persons
Limited exposure	2,000 – 3,000m	days to weeks	Acute mountain sickness (AMS) if the unacclimatized person sleeps at this altitude Beside AMS there is normally no risk to any person who does not have severe cardiopulmonary illness (HAPE is extremely rare at this altitude)
Expatriates	3,000 – 4,500m	years	high altitude pulmonary hypertension insufficiency of the right ventricle chronic mountain sickness (CMS, “Monge’s Disease”)
Altitude populations	>3,000m.	several generations	CMS (see above) Re-entry pulmonary oedema

Table 1: Characteristics of the types of exposure




Figure 2: pO2 and isobaric %O2 in relation to altitude. Graphs according to ICAO standard atmosphere as given by [12] (see also table 2). For fire protection oxygen concentration between 14.8 and 17% is used in most cases [13], [14].

Within the “extreme short exposure group” there is one small subgroup of subjects who are a special case: Persons who coach people, normally mountaineers, who pre-acclimatize for extreme altitude expeditions or employees who pre-acclimatize for work at high altitude. This is increasingly used, often in facilities which offer isobaric hypoxia. This may expose participants to an altitude of 5,300m or more. In most cases this exposure is limited for some minutes to up to half an hour. With the special advantage of isobaric hypoxia these persons can easily escape to normal atmospheric conditions at any time if they feel unwell. People with some pre-existing conditions may suffer from severe problems at this altitude while healthy persons should tolerate this exposure well: The duration of exposure is too short to develop acute mountain sickness (AMS) and it is also well below the limits of the period after which significant neurological risks may arise. In aviation medicine this period is called “time of useful consciousness” (table 2). Neither are relevant for this group.

Altitude	%O ₂ , isobaric conditions	Atmospheric pressure		pO ₂		Time of useful consciousness
		[mmHg]	[hPa]	[mmHg]	[hPa]	
[m]						
0	20,9	760.0	1013.2	158.8	211.7	No limitation
500	19,7	716.0	954.6	149.6	199.5	
1000	18,5	673.8	898.3	140.8	187.7	
1500	17,4	634.0	845.3	132.5	176.7	
2000	16,4	596.0	794.6	124.6	166.1	
2500	15,4	560.0	746.6	117.0	156.0	
3000	14,5	525.8	701.0	109.9	146.5	
3500	13,6	493.0	657.3	103.0	137.3	
4000	12,7	462.0	616.0	96.6	128.8	
4500	11,9	432.6	576.8	90.4	120.5	
5000	11,1	404.8	539.7	84.6	112.8	
5500	10,4	378.6	504.8	79.1	105.5	>30 min.
6000	9,7	353.6	471.4	73.9	98.5	
6500	9,1	330.0	440.0	69.0	92.0	
7000	8,5	307.8	410.4	64.3	87.7	3-5 min.
10500	5,0	183.0	244.0	38.2	50.9	ca. 1 min.
12900	3,4	123.5	164.7	25.8	34.4	15-30 sec.

Table 2: Atmospheric conditions (altitude, pressure, pO₂ and the corresponding % O₂ in isobaric hypoxia) according to ICAO standard atmosphere [12] and the time of useful consciousness for unacclimatized persons [15], [14]. There is no limitation of the time of useful consciousness up to 5000m or 11.1% O₂, respectively. For corresponding O₂% see also note on figure 2.

With increasing altitude the maximal workload decreases by 10% to 15% per 1,000m of altitude (beginning at 1,500m above sea level) with highly trained persons losing most power (figure 3). Since most work performed at altitude is of limited load on the human body (estimated 0.5 – 1.0 W/kg body weight) this effect is not limiting most of the previously described activities at altitude. During very demanding or exhausting work at >3,000m O₂ diffusion becomes a more and more limiting factor and persons doing intense work cannot stabilize their SaO₂ at a level which can be expected for the respective altitude. As a result their SaO₂ will decrease. Working conditions like

these are exclusively for healthy people, and even then the limited (decreased) workload must be taken into account when the required activity and the resources are planned (figure 3).




Figure 3: Decrease in maximal workload at altitude [16], [17], [18], [19], [20]. 1,500m: “threshold altitude”; 5,300m: limit of complete acclimatization / permanent stay

2.2 Limited Exposure

The typical exposure of the **Limited Exposure Group** is **2,000 to 3,000m** for some **days or weeks**. At some places these people will be exposed to even higher conditions of 4.000 – 5.000m, e.g. at Colorado or South America. Examples of locations for business men or normal travellers (not mountaineers) are given in figure 4. In contrast to the “Extreme Short Exposure Group” the combination of altitude and duration of exposure is potentially able to cause altitude disease (acute mountain sickness, AMS). An increase of the pulse rate for about 12-14% should be expected at 2,500-3,000m (e.g. [21], [13]). A comparable increase of minute ventilation and breathing frequency will occur [22]. Oxygen saturation will decrease only for about 6-8% of the saturation measured at sea level and be stable at 90-94% [23], [24]. A slight decrease of aerobic performance (without significant effect on performance at work) must be expected at 3,000m (figure 3), but no significant effect on psychomotoric functions [25], [26]. There are some data which indicate a slight impairment in the coordination of complex movements, especially when they have to be performed very quickly.




Figure 4: Examples for destinations of business men or normal travellers. Note: Some places are higher than the highest mountains in the European Alps

The main risk for the limited exposure group is the development of altitude diseases, most often AMS. High altitude pulmonary oedema (HAPE) is extremely rare at this altitude (1:4,000 nights at 3,000m [27]) as it normally occurs above 4,000m. A significantly increased risk exists for any person who has a rapid ascent profile, e.g. business men arriving by aircraft. If the arrival destination is around 4,500m the risk for HAPE with severe symptoms increases to 1:600 [27] (signs for HAPE in X-ray up to 31,7% [28]) and 30 – 57% will suffer from AMS [29], [30], [5], [31], [32]. On the other hand, these data indicate that some hours or even a night at 3,000m is relatively safe, at least for healthy people. Some people with pre-existing diseases may be at specific risk [1]. Nevertheless any strategy for occupational safety and health for the “limited exposure group” has to focus on AMS.

Note: In contrast to the belief of most Westerners many porters in the Himalayans are not true Sherpas anymore, and are instead immigrants or expatriate lowlanders! They may suffer from the same altitude-related risks as any other visitor [41].

Note: Persons who suffered from a stroke in their history, who got radiation therapy of neck or head, with large foramen ovale, or single pulmonary artery may be at increased risk when exposed to altitude, even if they do not have symptoms at low altitude (several case reports in literature, no systematic data). There are no data which support a general inability of these persons to be exposed to hypoxia or altitude, but they persons should be monitored closely during the first “test-“exposure, which should take place where it is easy to “escape”. Technical and medical surveyed rooms with isobaric hypoxia provide the optimum of safety for such testing.

2.3 Expatriates and Immigrants

Expatriates in this paper are defined as persons who migrate as lowlanders to high altitude regions, normally **above 3,000m**. Some of them are living higher than 4,500m. The duration may be for several months but is frequently for **years**. Normally this group will not suffer from acute altitude diseases (after the first few days at altitude) and they fully acclimatize.

The definition also includes immigrants. In many cases the employees are attended by their families. As a consequence, pregnancy may be a problem [9] and more often children of different ages will be exposed to high altitude. After a while they may suffer from specific diseases of the cardiopulmonary system: high altitude pulmonary hypertension (HAPH) (formerly called sub-acute mountain sickness) with acute bouts of high blood pressure in the pulmonary vessels [33] and high altitude heart disease. For details see [11].

After some months at high altitude adults may develop the symptoms of HAPH, caused by chronic pulmonary hypertension. These patients suffer from signs of insufficiency of the right ventricle (peripheral oedema, dyspnoea, cough, and angina pectoris). After years at altitude chronic mountain sickness (CMS, “Monge’s Disease”) may arise (headache, poor concentration, dizziness, reduced work capacity, cyanosis, clubbed / drumstick fingers, polycythaemia and high haemoglobin concentration) [34], [35], [36], [37], [38], [39], [40].

2.4 Altitude Populations

Altitude populations are defined as populations living for several **generations** (some for more than 30,000 years) at altitude **>3,000m**. These people show long-term and genetic adaptations. Most health problems are not altitude-related, but caused by socioeconomic factors (e.g. chronic bronchitis, Kangri carcinoma (skin cancer in cicatrice caused by multiple burning either by carrying a small oven heated with charcoal (kang) under the coat or by sleeping around a fire and burning the belly when the person turns around during sleep) etc.). While altitude-related health disorders are very uncommon in Tibetians and occupational medicine is not yet established in Ethiopia, South American highlanders – many of them working in mines – have been studied and are of special interest here. Some of them may develop CMS (see above). Another specific problem is re-entry pulmonary oedema where highlanders and also expatriates or immigrants who visit friends or relatives near sea level are of special risk when they return to their high altitude home after a stay of a week or two at low altitude. The symptoms are the same that for HAPE.

Note: In contrast to the belief of most Westerners, many porters in the Himalayans are not of true Sherpas anymore, and are immigrant or expatriate lowlanders! They may suffer from the same altitude-related risks as any other visitor [41]. Prefer to cooperate with agencies which follow the rules of the International Porters Protection Group (IPPG, see appendix 4).

3. Aspects of occupational safety and health

With the systematic structure given above, recommendations to provide specific advice and procedures for occupational medicine will be given here to facilitate personnel selection and offer health and safety guidance for employees going to altitude. An overview is given by the flowchart in appendix 2.

In contrast to any other occupational exposure where less exposure normally means less risk, employees at altitude benefit if they are exposed as often as possible to hypoxia because a partial adaptation can be expected (although scientific data about intermittent hypoxia are limited).

All persons working at altitude or in hypoxic conditions should be advised to drink enough to keep their fluid balance stable.

Note: In contrast to common thinking, it is not the heart which carries the greatest risk for people at altitude. While the myocardium is able to handle astonishing degrees of hypoxia the lung may be the limiting factor in the case of a pre-existing disease.

3.1 Extreme Short Exposure

As mentioned above the situation for this group may be best illustrated by the situation inside an aircraft's cabin (this excludes very high workload subjects and coaching of athletes who acclimatize to extreme altitudes, see below). In this well studied situation there are the following minimal requirements given by aviation medicine (values for sea level in adults) [42]:

- Vital capacity 3 l
- FEV1 70%
- SaO₂ 85%
- pO₂art. 70 mmHg

Haemoglobin should be >10g/dl and erythrocyte count >3 Million/ μ l [42]. Both are relative contraindications in the case of a chronically anaemic fully compensated patient who may be completely adapted.

For exposure up to 2,700m (equivalent to about 15.0-14.8% O₂) a check-up by Occupational Medicine – if any – should include the following topics:

- Person's history:
 - Any indicators for cardiopulmonary diseases or for significant limitations of maximum workload? (Sports?)
 - Any significant disease or operations in the last year?
 - Any problems during altitude sojourns in the past?
 - High or extreme workload at altitude expected?

If the person performs any aerobic endurance sport regularly and without problems, there is no doubt – without any other medical investigation – that there is no risk if this person is exposed within the conditions as described for the “extreme short exposure group” up to 2,700m or about 15.0-14.8% O₂.

- Laboratory or technical investigations are only required in cases which are not completely clear after checking the patient's history:

- Significant anaemia: blood status
 - Note: Exclude sickle cell anaemia if the person belongs to a group where this is endemic
- Pulmonary hypertension: echocardiography
- Pulmonary disease: spirometry and ergometry
- Cardiac disease: ergometry, stress echocardiography

If employees are exposed to equivalent altitudes of 2,700 – 3,800m as mentioned in chapter 2.1, the medical procedure should additionally include blood count, ergometry and spirometry (as $\dot{V}O_2\text{max}$ is the limiting factor at high altitude this may be combined as spiroergometry).

Employees who go to >3,800m should be checked as detailed above for 2,700-3,000m. Since hypoxic chambers are increasingly available (e.g. rooms or training centres for hypoxia training) these employees should be exposed to hypoxia once before their first departure to high altitude or before they enter hypoxic rooms for the first time. The exposure must take into account the kind, the expected duration, and especially the (equivalent) altitude (pO_2) of the exposure during work: For a task to be performed in a controlled environment (hypoxic rooms) for a limited duration, an acute exposure for 1-2 hours at the (equivalent) altitude where the work will be performed later should be sufficient. During exposure the person should be monitored for SaO_2 , pulse rate, and altitude-related symptoms by persons with adequate skills in altitude medicine. Persons with cardiopulmonary diseases >NYHA/CCS I, anaemic or pregnant persons should not be chosen for work at high altitude or hypoxia which corresponds to >3,800m [1], [43], [9].

For a normal work day it is not necessary to include additional pauses, if the real or equivalent altitude does not exceed 2,700m (rsp. 14.8% O_2 in isobaric conditions), as there is no risk for altitude-related disorders. If possible, and the work lasts all day in hypoxic rooms (e.g. fire protection), the person should be advised to leave the hypoxic area for a lunch break. If non-acclimatized persons are exposed to equivalent altitudes of 2,700 – 3,800m as mentioned above, they should be advised to make a pause in normoxia for at least 15 min. after every 2 hours of exposure. If they are exposed to >3,800m, this pause should be extended to 30 min. Non-acclimatized workers should avoid exposure to >4,500m.

Any employee should be advised to leave hypoxic areas if he / she does not feel well. To retreat to a normoxic area as soon as possible is the most important therapy in such a situation. As altitude-related diseases never start suddenly so any employee who heeds advice, which must be given to him prior to entering / ascending, will have plenty of time to escape from the hypoxia (see also table 2). It is not necessary to have a stand-by “rescue team” for this “extreme short duration group”.

If the symptoms recover completely after $\frac{1}{4}$ - $\frac{1}{2}$ hour, the person can re-enter, if necessary. If the subject does not recover within $\frac{1}{2}$ hour he / she should be advised to consult a physician who is a specialist in altitude medicine before the next exposure.

The employees should be advised to use a decongestant spray if they ascend quickly to altitude (hypobaric hypoxia) while suffering from upper airways infection (e.g. xylometazoline spray). Special care must be given to accompanying children in such situations [11].

Maximum workload need not be investigated for all employees: If the work which has to be performed at altitude is not very hard (e.g. paperwork (businessmen), supervi-

sion etc.) and the altitude does not exceed 2,700m, it is enough to know that the person is able to climb one flight of stairs or to walk 80-100m horizontal without shortage of breathing [43]. More detailed and according to NYHA / CCS stages [44], [45], [46] (for stages see also appendix 3):

- NYHA / CCS I (no symptoms): No limitations at altitude
- NYHA / CCS II (symptoms during moderate workload): No limitations at altitude for activities of minor workload
- NYHA III-IV (symptoms during minimal workload or at rest): Contraindication for altitude. Diagnoses and situations which are a permanent or temporary contraindication are given in table 2. We also refer to the recommendations given in [1].

Diagnosis	Period for which altitude / hypoxia should be avoided
Stroke	3 months
Myocardial infarction	6 weeks, if no complications 10 weeks (or more) in the case of complications (e.g. significant arrhythmia)
Aortocoronary bypass operation	2-3 weeks
PTCA	3 days
Stent	3-10 days
Pace maker implantation	After a check for correct function is o.k.
Acute broncho-pulmonary infections	No exposure until recovery
Asthma (stress induced)	No exposure, if not sufficiently treated

Table 3: Permanent or temporary contraindications for altitude sojourns of the “extreme short exposure group” as defined above (according to the recommendations for aircraft travellers) [42]

At altitudes which are normally visited for extreme short duration (1,500-3,000m) the maximal workload is only marginal impaired (fig.3). Therefore the recommendations given for demanding work (e.g. self containing breathing apparatuses) can be used without modification, if such work has to be done at 2,000-3,000m. Above that there is a 10% reduction per 1,000m above 1,500m which should be added to the minimal requirements, e.g.:

If a minimal requirement for a certain work at sea level is 2 W/kg body weight and this work shall be done at 4,000m, the estimation is as follows: $(4000-1500)/1000=2.5$; $2.5*10\%=25\%$; $2W/kg+25\% = 2.5 W/kg$. At sea level this employee must be able to provide at least 2.5 W/kg to be able to do the respective work at 4,000m.

Data about the workload necessary for several tasks at sea level are available from medical literature, e.g. [47]. An example for a protocol for a medical check for crew members of alpine helicopter rescue services is given as an example for employees at altitude exposed to high workload in appendix 1.

Persons who are exposed to extreme altitude e.g. while training expedition members must have a cardiopulmonary (especially pulmonary!) capacity well within in the normal range. This may include also well treated asthmatic patients. Above 4,500m em-

employees should be observed for safety by a “rescue person” who is outside the isobaric hypoxia but in permanent contact to the person(s) inside. For optimum supervision rooms for isobaric hypoxia should be constructed with a big window to the next room. Above 4,500m safety may be further increased if an oxygen bottle and a mask is available. Above 4,500m a specialist high altitude physician should be on call. Above 5,000m he/she should be on scene.

In contrast to others of the “extreme short exposure group” employees working in such conditions need to be educated about symptoms and management of acute hypoxia. Again: if they work regularly in such conditions (e.g. every day 2-3x for 15-60 min., or even more) a benefit by partial acclimatization can be expected (although data are scarce).

3.2 Limited Exposure

As this group goes to comparable altitude as the latter, the minimal requirements of the cardio-pulmonary system and the exclusion criteria (table 2) are the same. Special attention should be paid to persons who might be suspected for obstructive sleep apnoea, as this is often combined with pulmonary hypertension. The later will increase significantly at altitude. As mentioned in chapter 2 there is another risk for this group which cannot (yet) be predicted by any medical investigation: AMS. Here preventive occupational medicine includes detailed advice about the symptoms of AMS and how to treat them [48]. If possible discussion with the manager who is responsible for the employee and advice for an adequate ascent profile should be given. This may include an extra day on arrival and an overnight stay at intermediate altitude. If an immediate ascent and overnight stay >2,500m cannot be avoided prophylaxis with drugs may be considered (acetazolamide 125 mg bid, [48]).

It should be mentioned here that some facilities where people work at high altitude provide oxygen enriched rooms (e.g. some mines or telescopes in the Andes). Every additional 1% of oxygen corresponds to a “reduction” of 300m of altitude without increasing the risk of fire. With this technology some facilities provide an interior environment which corresponds to 3,000m, which is well tolerated.

If the person will be exposed without having the chance to escape from hypoxia (e.g. business trips to the Andes (South America)), an ascent profile according to international recommendations (e.g. [48]) should be an integral part of the strategy when the respective work is planned. If the place of work is situated higher than 3,800m the person should sleep at least one night under controlled conditions (hypobaric room at a training centre with physicians experienced in altitude medicine) at the respective equivalent altitude before his / hers first departure to such places. Depending on the final altitude this may include several nights in hypobaric conditions to realize an altitude profile and acclimatization according to the “Gold Standard” as recommended [48].

Any employee should be advised to contact an experienced high altitude physician immediately in the event that he / she does not feel well at high altitude and cannot descend. Painkillers, nifedipine, dexamethasone and acetazolamide with education and information on how to contact the physician to manage an emergency should be given to him / her before departure for any destination above 3,800m. Below 3,800m a standard painkiller (not aspirin!) should be sufficient.

2.3 Expatriates and Immigrants

For their travel to the destination this group should be informed about AMS as described in 2.2. Before they leave, they should get a more detailed medical examination, especially with electrocardiography and echocardiography included. This aims at two topics: i. to exclude pulmonary hypertension, and ii. to get initial values for a comparison with later controls. HAPH, the main risk for this group, causes a hypertrophy and dilatation of the right ventricle, a pulmonary hypertension, and sometimes a pericardial effusion. In many places where these expatriates work there will be no possibility for echocardiography. Then the follow up investigation may be performed during a holiday trip home. A control echocardiography should be performed at least once a year or if symptoms should arise. Note that accompanying children need special attention [11]. If any pathologies which indicate increased pulmonary pressure or right ventricle hypertrophy or dilatation are found at echocardiography the patients should be advised to leave their high altitude residency as soon as possible.

3.4 Altitude Populations

As mentioned above, altitude related problems are rare in this group. Employees descending from altitude for holidays or for other reasons for more than a week should be advised, as described in 2.2, to avoid re-entry HAPE. They should also be educated about First Aid if a HAPE should occur.

South American highlanders should be surveyed by occupational medicine for CMS. As a minimum these checks should include an analysis of haematocrit, haemoglobin, and erythrocyte cell count and should be performed annually. If symptoms arise which suggest CMS additional echocardiography should be performed, if possible.

References

1. Milledge J, Kupper T. Consensus Statement of the UIAA Medical Commission Vol.13: People with Pre-Existing Conditions Going to the Mountains. 2008 [cited 10.6.09]; Available from: www.theuiaa.org/act_medical.
2. Gunga HC, Kirsch K, Rucker L, Schobersberger W. Time course of erythropoietin, triiodothyronine, thyroxine, and thyroid-stimulating hormone at 2,315 m. *J Appl Physiol* 1994; 76(3): 1068-72.
3. Sakata S, Shimizu S, Kishi T, Hirai K, et al., Correlation between erythropoietin and lactate in humans during altitude exposure. *Jpn J Physiol* 2000; 50(2): 285-8.
4. Cottrell JJ, Altitude exposures during aircraft flight. *Flying higher*. *Chest* 1988; 93(1): 81-4.
5. Basnyat B, Acute mountain sickness in local pilgrims to a high altitude lake (4154 m) in Nepal. *J Wild Med* 1993; 4: 286-292.
6. Basnyat B, Murdock DR, High-altitude illnesses. *Lancet* 2003; 361: 1967-1974.
7. Rabold MB, Dexamethasone for prophylaxis and treatment of acute mountain sickness. *J Wilderness Med* 1992; 3(1): 54-60.
8. Hackett PH, Roach RC, High-altitude illness. *N Engl J Med* 2001; 345(2): 107-114.

9. Jean D, Leal C, Meijer H. Consensus Statement of the UIAA Medical Commission Vol.12: Women Going to Altitude. 2008 [cited 10.6.09]; Available from: www.theuiaa.org/act_medical.
10. Jean D, Leal C, Kriemler S, Meijer H, et al., Medical recommendations for women going to altitude. *High Alt Med Biol* 2005; 6(1): 22-31.
11. Meijer HJ, Jean D. Consensus Statement of the UIAA Medical Commission Vol.9: Children at Altitude. 2008 [cited 10.6.09]; Available from: www.theuiaa.org/act_medical.
12. Ruff S, Strughold H, Grundriss der Luftfahrtmedizin. 2. Aufl. ed. Leipzig, Johann Ambrosius Barth, 1944. 249.
13. Muller B, Die gesamte Luftfahrt- und Raumflugmedizin. Düsseldorf, Droste Verlag, 1967.
14. Ernsting J, King P, Aviation Medicine. 2nd ed. ed. Oxford, Butterworth-Heinemann Ltd, 1994.
15. Amsler HA, Flugmedizin für zivile Besatzungen. Bern, Verlag Eidgenössisches Luftamt, 1971. 135.
16. Kupper T, [Workload and professional requirements for alpine rescue]. Professoral thesis at Aachen Technical University / Germany, 2006 (english publication in preparation), in Dept. of Aerospace Medicine. 2006, Aachen Technical University: Aachen.
17. West JB, Limiting factors for exercise at extreme altitudes. *Clin Physiol* 1990; 10(3): 265-72.
18. Buskirk ER, Kollias J, Picon Reategui E, Physiology and performance of track athletes at various altitudes in the United States and Peru, in The international symposium on the effects of altitude on physical performance, R.F. Goddard, Editor. 1966, The Athletic Institute: Chicago.
19. Jackson CG, Sharkey BJ, Altitude, training and human performance. *Sports Med* 1988; 6(5): 279-84.
20. Buskirk ER, Kollias J, Akers RF, Prokop EK, et al., Maximal performance at altitude and on return from altitude in conditioned runners. *J Appl Physiol* 1967; 23: 259-267.
21. Glaisher J, Notes of effects experienced during recent balloon ascents. *Lancet* 1862; 2: 559-560.
22. Rupwate RU, Chitaley M, Kamat SR, Cardiopulmonary functional changes in acute acclimatization to high altitude in mountaineers. *Eur J Epidemiol* 1990; 6(3): 266-72.
23. Horii Xea, *J.Wild.Med.* 1994; 5: 447-450.
24. Cottrell JJ, Lebovitz BL, Fennell RG, Kohn GM, Inflight arterial saturation: continuous monitoring by pulse oximetry. *Aviat Space Environ Med* 1995; 66(2): 126-30.
25. Waanders R, Riedmann G, Short term impairment in cognitive functioning after a rapid ascent to altitude of 4,559 meters (Abstract). *Eur J Neurisci (Suppl)* 1994; 7: 217.
26. Tune GS, Psychological effects of hypoxia: Review of certain literature from the period 1950 to 1963. *Percept Mot Skills* 1964; 19: 551-562.
27. Hochstrasser J, Nanzer A, Oelz O, [Altitude edema in the Swiss Alps. Observations on the incidence and clinical course in 50 patients 1980-1984]. *Schweiz Med Wochenschr* 1986; 116(26): 866-73.
28. Bircher HP, Eichenberger U, Maggiorini M, Oelz O, et al., Relationship of mountain sickness to physical fitness and exercise intensity during ascent. *J Wild Med* 1994; 5(4): 302-311.
29. Basnyat B, Lemaster J, Litch JA, Everest or bust: a cross sectional, epidemiological study of acute mountain sickness at 4243 meters in the Himalayas. *Aviat Space Environ Med* 1999; 70(9): 867-73.
30. Hackett PH, Rennie D, Rales, peripheral edema, retinal hemorrhage and acute mountain sickness. *Am J Med* 1979; 67(2): 214-8.
31. Maggiorini M, Buhler B, Walter M, Oelz O, Prevalence of acute mountain sickness in the Swiss Alps. *Bmj* 1990; 301(6756): 853-5.
32. Schneider M, Bernasch D, Weymann J, Holle R, et al., Acute mountain sickness: influence of susceptibility, preexposure, and ascent rate. *Med Sci Sports Exerc* 2002; 34(12): 1886-91.

33. Sui GJ, Liu YH, Cheng XS, Anand IS, et al., Subacute infantile mountain sickness. *J Pathol* 1988; 155(2): 161-70.
34. Arregui A, Leon-Velarde F, Cabrera J, Paredes S, et al., Migraine, polycythemia and chronic mountain sickness. *Cephalalgia* 1994; 14(5): 339-41.
35. Bernardi L, Roach RC, Keyl C, Spicuzza L, et al., Ventilation, autonomic function, sleep and erythropoietin. Chronic mountain sickness of Andean natives. *Adv Exp Med Biol* 2003; 543: 161-75.
36. Curran LS, Zhuang J, Sun SF, Moore LG, Ventilation and hypoxic ventilatory responsiveness in Chinese-Tibetan residents at 3,658 m. *J Appl Physiol* 1997; 83(6): 2098-104.
37. Ge RL Helun G, Current concept of chronic mountain sickness: pulmonary hypertension-related high-altitude heart disease. *Wilderness Environ Med* 2001; 12(3): 190-4.
38. Leon-Velarde F, Reeves JT, International consensus group on chronic mountain sickness. *Adv Exp Med Biol* 1999; 474: 351-3.
39. Monge CC, Arregui A, Leon-Velarde F, Pathophysiology and epidemiology of chronic mountain sickness. *Int J Sports Med* 1992; 13 Suppl 1: S79-81.
40. Moore LG, Niermeyer S, Zamudio S, Human adaptation to high altitude: regional and life-cycle perspectives. *Am J Phys Anthropol* 1998; Suppl 27: 25-64.
41. Basnyat B, Litch JA, Medical problems of porters and trekkers in the Nepal Himalaya. *Wilderness Environ Med* 1997; 8(2): 78-81.
42. Siedenburg J, *Kompodium Reisemedizin und Flugmedizin*. 6. Aufl. ed. Norderstedt, BoD - Books on Demand, 2009.
43. Kupper T, [Non-traumatic aspects of sport climbing]. *Wien Med Wochenschr* 2005; 155(7-8): 163-70.
44. N.N., The Criteria Committee of the New York Heart Association. *Diseases of the Heart and Blood Vessels: Nomenclature and Criteria for Diagnosis*, in *The Criteria Committee of the New York Heart Association. Diseases of the Heart and Blood Vessels: Nomenclature and Criteria for Diagnosis*, N.N., Editor. 1928, Little Brown: Boston, Mass.
45. N.N., The Criteria Committee of the New York Heart Association. *Nomenclature and Criteria for Diagnosis of Diseases of the Heart and Great Vessels*, in *The Criteria Committee of the New York Heart Association. Nomenclature and Criteria for Diagnosis of Diseases of the Heart and Great Vessels*, N.N., Editor. 1994, Little, Brown & Co: Boston, Mass. p. 253-256.
46. Miller-Davis C, Marden S, Leidy NK, The New York Heart Association Classes and functional status: what are we really measuring? *Heart Lung* 2006; 35(4): 217-24.
47. Spitzer H, Hettlinger T, Kaminski G, *Tafeln für den Energieumsatz bei körperlicher Arbeit*. Berlin, Köln, Beuth Verlag, 1982.
48. Kupper T, Gieseler U, Angelini C, Hillebrandt D, et al. Consensus Statement of the UIAA Medical Commission Vol.2: Emergency Field Management of Acute Mountain Sickness, High Altitude Pulmonary Oedema, and High Altitude Cerebral Oedema. 2008 [cited 2008 11.1.09]; Available from: www.theuiaa.org/act_medical.

Members of UIAA MedCom

C. Angelini (Italy), B. Basnyat (Nepal), J. Bogg (Sweden), A.R. Chioconi (Argentina), S. Ferrandis (Spain), U. Gieseler (Germany), U. Hefti (Switzerland), D. Hillebrandt (U.K.), J. Holmgren (Sweden), M. Horii (Japan), D. Jean (France), A. Koukoutsis (Greece), J. Kubalova (Czech Republic), T. Küpper (Germany), H. Meijer (Netherlands), J. Milledge (U.K.), A. Morrison (U.K.), H. Mosaedian (Iran), S. Omori (Japan), I. Rotman (Czech Republic), V. Schöffl (Germany), J. Shahbazi (Iran), J. Windsor (U.K.)

History of this recommendation paper

The version presented here was approved by written consent in lieu of a live meeting in August 2009

Appendix 1: Occupational Medicine Check for Crew Members of Alpine Helicopter Rescue Services [16]

1. First check (before start of work)

1.1 Patient's history

- General history
- Specific history (aviation medicine and altitude medicine)
 - Problems during or after flight (which ones?, when?, reproducible?, at which altitude?, how long after arrival at altitude?, other symptoms?)
 - Problems linked to an altitude sojourn, especially shortness of breath at altitude, dyspnoea or reduced fitness (when?, reproducible?, at which altitude?, when after arrival at altitude?, other symptoms?); AMS, HAPE or HACE in history?
- Vaccinations
 - Previous or actual infections or infectious diseases
- Work history
- Actual symptoms
- Attention should be paid to:
 - Immunological diseases or diseases (or therapies) which significantly impair the immunological system
 - Circulatory disturbance (hypertension, hypotension with dizziness)
 - Coronary heart disease / angina pectoris, history of myocardial infarction, arrhythmia, cardiac insufficiency
 - Cerebral trauma
 - Diabetes mellitus (especially Type I A)
 - Nephropathy
 - Dermatological diseases which facilitate the invasion of pathogenic germs or which cause an increased sensitivity to UV radiation
 - Neurological diseases
 - Psychiatric diseases
 - Use of drugs or luxury food which may have sedative side effects, alcohol or hypnotic drugs
 - Ophthalmological diseases or trauma which impair the visibility
 - Diseases or trauma of the middle or inner ear
 - Sensitivity to kinetosis
 - Psychic resilience (acrophobia, free of dizziness, fear of flying)

1.2 Medical check with special regard to the kind of work

- Clinical check-up
 - Screening test „Noise“
 - Otoscopy
 - Audiometry (air conduction) at 1-6 kHz
 - Laboratory investigations
 - Urine screening (dipstick)
 - ESR
 - Blood count (haemoglobine, erythrocytes, leucocytes, haematocrite)
 - GGT, GOT, GPT
 - Creatinine
 - Serum glucose concentration (where appropriate: oral glucose stress test, Hb_{A1c})
 - Anti-HBc or (vaccinated persons) Anti-HBs quantitatively
 - If Anti-HBc positive: check HBs-Ag and Anti-HBs quantitatively
 - If HBs-Ag positive: check HBe-Ag and Anti-HBe
 - Eyes
 - Measurement of visus (proximity and distance)
 - Stereoscopic vision
 - Colour vision
 - Field of vision
 - Mesopic vision sensibility to glare
-

- Thorax X-ray
- Performance
 - Spirometry
 - Ergometry
- Balance
 - Romberg test (1 min.)
 - Fukuda test (1 min.)

Remarks:

- Audiometry is not necessary if the respective data are available from a check which was performed <6 months before. If any pathologies are verified an immediate extended check of the ears is indicated.
- Thorax X-ray is not necessary if an X-ray which was performed <2 years ago is available and when no significant diagnosis is expected.

1.3 Additional investigations which may be performed

- Cranio-Corpography with photooptical recording
- Spiroergometry

1.4 Criteria of aviation medicine and occupational medicine

1.4.1 Permanent unable for work at altitude or in hypoxia

- Persons with one or several of the following neurological, otological, or psychiatric situation::
 - Impaired consciousness or seizure disorder independent from the cause
 - Neurological diseases (central or peripheral) with significant functional impairment, especially after cerebral trauma, impairment of the cerebral perfusion, organic diseases of the brain or the spinal cord, and peripheral neuropathies independent from the cause
 - Disturbance of equilibrium with
 - Lateral oscillations in the **Fukuda** stepping test >20 cm or lateral deviance in the Romberg test >80° to the right or >70° to the left side, or
 - Longitudinal oscillations in the Romberg test of >12 cm or lateral oscillations >10 cm, if these did not disappear after the observation periods given in 1.4.2
 - Chronic attacks of dizziness with significant vestibulo-okular or retino-okular disturbance of movement (to be proved in electromyostagmy)
 - Permanent medication with drugs which may have sedative side effects.
 - Any kind of drug or stimulant abuse, also if in the person's history
 - Any psychic disease, also if in the case of a significant amelioration a relapse or decompensation may have a significant influence on occupational safety (especially diseases which may impair stress management or assessment of risk)
 - An individually increased risk for hearing loss by noise, e.g.:
 - Significant hearing loss in bone conduction of at least one ear and at least 1 test frequency between 1 and 6 kHz
 - Any vestibular dizziness or Morbus Menière
 - Any significant diseases of the inner ear in the person's history (e.g. acute hearing loss)
 - Hardness of hearing of the inner ear or by Nervus cochlearis, or after cerebral trauma
 - Otosclerosis operation in the person's history
 - Persons who do not tolerate ear protection, headsets, or helmets with integrated communication systems because of any disease of the ear canal or the auricle (e.g. ekzema which is resistant to therapy)
 - Persons who are unable to perform pressurization of the ear or the sinuses because of a chronic disease
 - Persons with chronic disease of the external parts of the eyes
- Persons, who apply to one or several of the following aspects of internal medicine or performance:
 - Any disease of the cardio-circulatory system which causes a permanent impairment of performance or the ability for regulation and which in the long run limit the individual performance to a PWC170 of <3,0 W/kg body weight [16] (especially coronary heart

- disease, cardiac insufficiency, pulmonary hypertension, or severe arterial hypertension)
- Any disease of the respiratory system which causes a permanent impairment of performance or the ability for regulation and which in the long run limit the individual performance to a PWC170 of <3,0 W/kg body weight [16] (especially chronic-obstructive pulmonary disease (COPD), diseases of the pulmonary tissue which cause restrictive breathing disturbance, any kind of disturbances of gas exchange)
 - Any change of the red blood cells which causes a permanent impairment of performance to a PWC170 of <3,0 W/kg body weight [16] (e.g. chronic anaemia independent from its cause). Any disease of the red blood cells or anomalies of haemoglobine which may decompensate acute in hypoxia (e.g. sickle cell anaemia)
 - Any disease of the skin or the blood vessels which impairs the peripheral perfusion (risk for frostbite!) or persons with frostbite >I° in their history.
 - Significant diseases of the kidney or urinary system (e.g. insufficiency III° or IV° with a glomerular filtration rate of 30 – 59 ml/min [6])
 - Rheumatic diseases with the risk of worsening if exposed to cold climate and all severe stages of rheumatic diseases
 - Any metabolic disease which causes a permanent impairment of performance, alertness, coordination or balance (e.g. diseases of the thyroid gland, parathyroid, adrenal glands)
 - Medically treated diabetes, especially if the patient tends to hypoglycaemia
 - Persons with intolerance reactions to cold exposure (e.g. cold agglutinins, cold urticaria, or cold haemoglobinuria)
 - Permanent reduction of the muscular strength, the flexibility, or with the loss of any limb if the impairment causes a permanent degree that the realization of the task cannot be expected.
 - Untreated sleep disturbances (sleep apnoea) if they cause a significant impairment of alertness or pulmonary hypertension
 - Persons with reduced immunological competence like:
 - Chronic diseases or long-ranging infections (e.g. HIV), which cause a significant impairment of the immune defence
 - Therapy with immune suppressive drugs, cytostatic drugs, or ionising radiation
 - Chronic (therapy resistant) eczema of the hands, which impair the protecting effect of the skin against infectious agents
 - Persons with chronic skin diseases, which cause an increased sensibility against UV radiation

1.4.2 Temporary concerns

Persons with impairment of health listed in 1.4.1, if a complete or sufficient recovery can be expected, e.g.:

- Persons with a fitness of less than 3.0 W/kg body weight (PWC170) [16]: create training schedule, next medical check after 6-12 months depending on the performance and the compliance of the person
 - Persons with anaemia <11,0 g/dl, if a recovery to normal values can be expected
 - Temporary reduced immune defence (e.g. during temporary corticoid medication with high dosages or during severe acute infectious diseases)
 - Temporary reduced ability for pressurization (e.g. during acute infections of the upper respiratory system)
 - Persons with a body-mass-index >28 or a body weight >90 kg (the limit of the helicopter winch includes the weight of the rescuer, of the patient and of the equipment!)
 - Acute eczema of the hands or large-scale abrasion, which impair the protective effect of the skin against infectious agents for a while
 - Persons who exceed the limits of Romberg test and Fukuda test as given above. Annual re-check to test for possible recovery. After the 4th year any amelioration cannot be expected. The latter situation causes permanent concerns and the person has to quit the job.
 - Persons with temporary disease, which make the use of hearing protection, headsets, or helmets with integrated communication devices impossible for a while (e.g. injuries of the external ear, acute disease of the auditory canal or of the auricle)
-

- Persons who temporarily have to take drugs which may increase sensitivity to UV radiation

1.4.3 No concerns if specific preconditions are realized

- Immune defence: If the disturbance is less significant, a careful consideration should be done, whether a further activity may be accepted (define specific preventive procedures, where appropriate). Reduced period until medical re-check recommended.
- Migraine (specific need for advice, prevention of attacks by drugs, if appropriate)

1.4.4 No concerns at all

Any other persons, as far as there are no restrictions or a ban by laws (e.g. pregnant women, youths)

2 Re-checks

2.1 Periods until re-check

2.1.1 First re-check: before the end of the 6th month after beginning of the work


2.1.2 Further re-checks: before a 12-months-period ends

2.1.3 Early re-check: If any disease with a duration of >4 weeks or any disease which may impair significantly the performance of the cardiopulmonary or the peripheral muscular system, also if the employee assumes a correlation between his symptoms or disease and the work.

3 Medical check after the employee has finished the job

On the eve of 6 months after finishing the job a serological check for Hepatitis B/C and HIV is recommended.

Appendix 2: Flowchart how to manage occupational health and safety for work in hypoxia



Appendix 3: New York Heart Association Heart Failure Classification (revised version 1994) [45]

Functional Capacity	Objective Assessment
Class I. Patients with cardiac disease but without resulting limitation of physical activity. Ordinary physical activity does not cause undue fatigue, palpitation, dyspnoea, or anginal pain.	A. No objective evidence of cardiovascular disease.
Class II. Patients with cardiac disease resulting in slight limitation of physical activity. They are comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnoea, or anginal pain.	B. Objective evidence of minimal cardiovascular disease.
Class III. Patients with cardiac disease resulting in marked limitation of physical activity. They are comfortable at rest. Less than ordinary activity causes fatigue, palpitations, dyspnoea, or anginal pain.	C. Objective evidence of moderately severe cardiovascular disease.
Class IV. Patients with cardiac disease resulting in inability to carry on any physical activity without discomfort. Symptoms of heart failure or the anginal syndrome may be present even at rest. If any physical activity is undertaken, discomfort is increased.	D. Objective evidence of severe cardiovascular disease

Appendix 4: Guidelines of the International Porter Protection Group (IPPG)

1. Trekking ethics

1. Clothing that is appropriate for season and altitudes encountered must be provided to porters for protection from cold, rain and snow. This may mean: wind-proof jacket and trousers, fleece jacket, long johns, suitable footwear (leather boots in snow), socks, hat, gloves and sunglasses.
2. Above the tree line porters should have a dedicated shelter, either a room in a lodge or a tent (the trekkers' mess tent is no good as it is not available till late evening), a sleeping pad and a blanket (or sleeping bag). They should be provided with food and warm drinks, or cooking equipment and fuel.
3. Porters should be provided with the same standard of medical care as you would expect for yourself, and life insurance.
4. Porters should not be paid off because of illness/injury without the leader or the trekkers assessing their condition carefully. The person in charge of the porters (sirdar) must let their trek leader or the trekkers know if a sick porter is about to be paid off. Failure to do this has resulted in many deaths. Sick/injured porters should never be sent down alone, but with someone who speaks their language and understands their problem, along with a letter describing their complaint. Sufficient funds should be provided to cover cost of rescue and treatment.
5. No porter should be asked to carry a load that is too heavy for their physical abilities (maximum: 20 kg on Kilimanjaro, 25 kg in Peru and Pakistan, 30 kg in Nepal). Weight limits may need to be adjusted for altitude, trail and weather conditions; experience is needed to make this decision.

2. Questions to ask trekking companies:

1. Does the company you are thinking of trekking with follow IPPG's five guidelines on porter safety?
2. What is their policy on equipment and health care for porters?
3. What do they do to ensure the trekking staff is properly trained to look after porters' welfare?
4. What is their policy on training and monitoring porter care by its ground operator in Nepal?
5. Do they ask about treatment of porters in their post trek questionnaire to clients?

From: www.ippg.net, accession date Aug. 3rd, 2008